

5. Dushianthan A, Knight M, Russell P, Grocott MP. Goal-directed haemodynamic therapy (GDHT) in surgical patients: systematic review and meta-analysis of the impact of GDHT on post-operative pulmonary complications. *Perioperative Medicine*. 2020;9(1). <https://doi.org/10.1186/s13741-020-00161-5>.
6. Giglio M, Biancofiore G, Corriero A, Romagnoli S, Tritapepe L, Brienza N, et al. Perioperative goal-directed therapy and postoperative complications in different kind of surgical procedures: an updated meta-analysis. *Journal of Anesthesia Analgesia and Critical Care*. 2021;1(1). <https://doi.org/10.1186/s44158-021-00026-3>.
7. Talmor D, Kelly B. How to better identify patients at high risk of postoperative complications? *Current Opinion in Critical Care*. 2017;23(5):417-423. <https://doi.org/10.1097/mcc.0000000000000445>.
8. POSE-Study group. Peri-interventional outcome study in the elderly in Europe. *European Journal of Anaesthesiology*. 2021;39(3):198-209. <https://doi.org/10.1097/eja.0000000000001639>.
9. Cecconi M, De Backer D, Antonelli M, Beale R, Bakker J, Hofer C, et al. Consensus on circulatory shock and hemodynamic monitoring. Task force of the European Society of Intensive Care Medicine. *Intensive Care Medicine*. 2014;40(12):1795-1815. <https://doi.org/10.1007/s00134-014-3525-z>.
10. ESICM Consensus Conference on shock diagnosis and haemodynamic monitoring in the ICU. The co-chairs are Michel Chew (University of Lund, Sweden) and Xavier Monnet (University of Paris Saclay, France).
11. Swan HJC, Ganz W, Forrester J, Marcus H, Diamond G, Chonette D. Catheterization of the Heart in Man with Use of a Flow-Directed Balloon-Tipped Catheter. *New England Journal of Medicine*. 1970;283(9):447-451. <https://doi.org/10.1056/nejm197008272830902>.
12. Gershengorn HB, Wunsch H. Understanding changes in established practice. *Critical Care Medicine*. 2013;41(12):2667-2676. <https://doi.org/10.1097/ccm.0b013e318298a41e>.
13. Robba C, Wong A, Poole D, Tayar AA, Arntfield RT, Chew MS, et al. Basic ultrasound head-to-toe skills for intensivists in the general and neuro intensive care unit population: consensus and expert recommendations of the European Society of Intensive Care Medicine. *Intensive Care Medicine*. 2021;47(12):1347-1367. <https://doi.org/10.1007/s00134-021-06486-z>.
14. Perkins GD, Gräsner J, Semeraro F, Olasveengen T, Soar J, Lott C, et al. European Resuscitation Council Guidelines 2021: Executive summary. *Resuscitation*. 2021;161:1-60. <https://doi.org/10.1016/j.resuscitation.2021.02.003>.
15. Stewart GN. Researches on the Circulation Time in Organs and on the Influences which affect it. *Journal of Physiology*. 1893;15(1-2):1-89. <https://doi.org/10.1113/jphysiol.1893.sp000462>.
16. Hamilton WF, Moore JW, Kinsman JM, Spurling RG. Studies on the circulation. IV. Further analysis of the injection method, and of changes in hemodynamics under physiological and pathological conditions. *Am J Physiol*. 1932;99:534-51.
17. Jain M, Canham M, Upadhyay D, Corbridge T. Variability in interventions with pulmonary artery catheter data. *Intensive Care Medicine*. 2003;29(11):2059-2062. <https://doi.org/10.1007/s00134-003-1924-7>.
18. Nishikawa T, Dohi S. Errors in the measurement of cardiac output by thermodilution. *Canadian Journal of Anaesthesia/Canadian Journal of Anesthesia*. 1993;40(2):142-153. <https://doi.org/10.1007/bf03011312>.
19. Galluccio ST, Chapman MJ, Finnis ME. Femoral-radial arterial pressure gradients in critically ill patients. *Critical Care and Resuscitation*. 2009;11(1):34-38. [https://doi.org/10.1016/s1441-2772\(23\)01830-6](https://doi.org/10.1016/s1441-2772(23)01830-6).
20. Goedje O, Hoeke K, Lichtwarck-Aschoff M, Faltchauser A, Lamm P, Reichart B. Continuous cardiac output by femoral arterial thermodilution calibrated pulse contour analysis: Comparison with pulmonary arterial thermodilution. *Critical Care Medicine*. 1999;27(11):2407-2412. <https://doi.org/10.1097/00003246-199911000-00014>.
21. Marx G, Schuerholz T, Sumpelmann R, Simon T, Leuwer M. Comparison of cardiac output measurements by arterial trans-cardiopulmonary and pulmonary arterial thermodilution with direct Fick in septic shock. *European Journal of Anaesthesiology*. 2005;22(2):129-134. <https://doi.org/10.1017/s0265021505000244>.
22. Della Rocca G, Costa MG, Coccia C, Pompei L, Di Marco P, Vilardi V, et al. Cardiac output monitoring: aortic transpulmonary thermodilution and pulse contour analysis agree with standard thermodilution methods in patients undergoing lung transplantation. *Canadian Journal of Anesthesia*. 2003;50(7):707-711. <https://doi.org/10.1007/bf03018714>.
23. Sangkum L, Liu GL, Yu L, Yan H, Kaye AD, Liu H. Minimally invasive or noninvasive cardiac output measurement: an update. *Journal of Anesthesia*. 2016;30(3):461-480. <https://doi.org/10.1007/s00540-016-2154-9>.
24. Kushimoto S, Taira Y, Kitazawa Y, Okuchi K, Sakamoto T, Ishikura H, et al. The clinical usefulness of extravascular lung water and pulmonary vascular permeability index to diagnose and characterize pulmonary edema: a prospective multicenter study on the quantitative differential diagnostic definition for acute lung injury/acute respiratory distress syndrome. *Critical Care*. 2012;16(6). <https://doi.org/10.1186/cc11898>.
25. Gavelli F, Shi R, Teboul F, Azzolina D, Mercado P, Jozwiak M, et al. Extravascular lung water levels are associated with mortality: a systematic review and meta-analysis. *Critical Care*. 2022;26(1). <https://doi.org/10.1186/s13054-022-04061-6>.
26. Shi R, Ayed S, Beuzelin M, Persichini R, Legouge M, Vita N, et al. Incidence and risk factors of weaning-induced pulmonary oedema: results from a multicentre, observational study. *Critical Care*. 2025;29(1). <https://doi.org/10.1186/s13054-025-05350-6>.
27. López-Herce J, Bustinza A, Sancho L, Mencía S, Carrillo A, Moral R, et al. Cardiac output and blood volume parameters using femoral arterial thermodilution. *Pediatrics International*. 2009;51(1):59-65. <https://doi.org/10.1111/j.1442-200x.2008.02654.x>.
28. Saugel B, Kouz K, Meidert AS, Schulte-Uentrop L, Romagnoli S. How to measure blood pressure using an arterial catheter: a systematic 5-step approach. *Critical Care*. 2020;24(1). <https://doi.org/10.1186/s13054-020-02859-w>.
29. Foti L, Michard F, Villa G, Ricci Z, Romagnoli S. The impact of arterial pressure waveform underdamping and resonance filters on cardiac output measurements with pulse wave analysis. *British Journal of Anaesthesia*. 2022;129(1):e6-e8. <https://doi.org/10.1016/j.bja.2022.03.024>.
30. Sagawa K. Translation of Otto Frank's paper "Die Grundform des arteriellen Pulses" zeitschrift für biologie 37:483-526 (1899). *Journal of Molecular and Cellular Cardiology*. 1990;22(3):253-254. [https://doi.org/10.1016/0022-2828\(90\)91459-k](https://doi.org/10.1016/0022-2828(90)91459-k).
31. Latham RD, Westerhof N, Sipkema P, Rubal BJ, Reuderink P, Murgo JP. Regional wave travel and reflections along the human aorta: a study with six simultaneous micromanometric pressures. *Circulation*. 1985;72(6):1257-1269. <https://doi.org/10.1161/01.cir.72.6.1257>.
32. Lee AJ, Cohn JH, Ranasinghe JS. Cardiac output assessed by invasive and minimally invasive techniques. *Anesthesiology Research and Practice*. 2011;1-17. <https://doi.org/10.1155/2011/475151>.
33. Saugel B, Kouz K, Scheeren TW, Greiwe G, Hoppe P, Romagnoli S, et al. Cardiac output estimation using pulse wave analysis – physiology, algorithms, and technologies: a narrative review. *British Journal of Anaesthesia*. 2021;126(1):67-76. <https://doi.org/10.1016/j.bja.2020.09.049>.
34. Mohammadi I, Firouzabadi SR, Hosseinpour M, Akhlaghpasand M, Hajikarimloo B, Tavanaei R, et al. Predictive ability of hypotension prediction index and machine learning methods in intraoperative hypotension: a systematic review and meta-analysis. *Journal of Translational Medicine*. 2024;22(1). <https://doi.org/10.1186/s12967-024-05481-4>.
35. an Der Ven WH, Terwindt LE, Risvanoglu N, le ELK, Wijnberge M, Veelo DP, et al. Performance of a machine-learning algorithm to predict hypotension in mechanically ventilated patients with COVID-19 admitted to the intensive care unit: a cohort study. *Journal of Clinical Monitoring and Computing*. 2021;36(5):1397-1405. <https://doi.org/10.1007/s10877-021-00778-x>.
36. Barile L, Landoni G, Pieri M, Ruggeri L, Maj G, Neto C, et al. Cardiac index assessment by the Pressure Recording Analytic Method in critically ill unstable patients after cardiac surgery. *Journal of Cardiothoracic and Vascular Anesthesia*. 2013;27(6):1108-1113. <https://doi.org/10.1053/j.jvca.2013.02.016>.
37. Joosten A, Desebbe O, Suehiro K, Murphy L, Essiet M, Alexander B, et al. Accuracy and precision of non-invasive cardiac output monitoring devices in perioperative medicine: a systematic review and meta-analysis. *British Journal of Anaesthesia*. 2016;118(3):298-310. <https://doi.org/10.1093/bja/aew461>.
38. Lamarche-Fontaneto R, Oud L, Howell KD, Ganerwal SA, Manek GU, Schleicher MC, et al. Cardiac output monitors in septic shock: do they deliver what matters? A systematic review and meta-analysis. *Critical Care*. 2025;29(1). <https://doi.org/10.1186/s13054-025-05547-9>.
39. Kobe J, Mishra N, Arya VK, Al-Moustadi W, Nates W, Kumar B. Cardiac output monitoring: Technology and choice. *Annals of Cardiac Anaesthesia/Annals of Cardiac Anaesthesia*. 2019;22(1):6-17. https://doi.org/10.4103/aca.aca_41_18.
40. Lefrant JY, Bruelle P, Aya AGM, Saïssi G, Dauzat M, De La Coussaye JE, et al. Training is required to improve the reliability of esophageal doppler to measure cardiac output in critically ill patients. *Intensive Care Medicine*. 1998;24(4):347-352. <https://doi.org/10.1007/s001340050578>.
41. Penáz J, Voigt A, Teichmann W. Beitrag zur fortlaufenden indirekten Blutdruckmessung [Contribution to the continuous indirect blood pressure measurement]. *Z Gesamte Inn Med*. 1976 Dec 15;31(24):1030-3. German. PMID: 1020409.
42. Maheshwari K, Khanna S, Bajracharya GR, Makarova N, Riter Q, Raza S, et al. A randomized trial of continuous noninvasive blood pressure monitoring during noncardiac surgery. *Anesthesia and Analgesia*. 2018;127(2):424-431. <https://doi.org/10.1213/ane.0000000000003482>.
43. Wagner JY, Grond J, Fortin J, Negulescu I, Schöfthaler M, Saugel B. Continuous noninvasive cardiac output determination using the CNAP system: evaluation of a cardiac output algorithm for the analysis of volume clamp method-derived pulse contour. *Journal of Clinical Monitoring and Computing*. 2015;30(4):487-493. <https://doi.org/10.1007/s10877-015-9744-1>.
44. Walsh M, Devereaux PJ, Garg AX, Kurz A, Turan A, Rodseth RN, et al. Relationship between Intraoperative Mean Arterial Pressure and Clinical Outcomes after Noncardiac Surgery. *Anesthesiology*. 2013;119(3):507-515. <https://doi.org/10.1097/aln.0b013e3182a10e26>.
45. Wesslink E, Kappen T, Torn H, Slooter A, Van Klei W. Intraoperative hypotension and the risk of postoperative adverse outcomes: a systematic review. *British Journal of Anaesthesia*. 2018;121(4):706-721. <https://doi.org/10.1016/j.bja.2018.04.036>.
46. Maleczek M, Laxar D, Geroldinger A, Gleiss A, Lichtenegger P, Kimberger O. Definition of clinically relevant intraoperative hypotension: A data-driven approach. *PLoS ONE*. 2024;19(11):e0312966. <https://doi.org/10.1371/journal.pone.0312966>.
47. Gregory A, Stapelfeldt WH, Khanna AK, Smischney NJ, Boero IJ, Chen Q, et al. Intraoperative hypotension is associated with adverse clinical outcomes after noncardiac surgery. *An-*