

## REFERENCES

1. Vivien B, Paqueron X, Le Cosquer P, Langeron O, Coriat P, Riou B. Detection of brain death onset using the bispectral index in severely comatose patients. *Intensive Care Med.* 2002;28(4):419-425. doi:10.1007/s00134-002-1219-4.
2. Escudero D, Otero J, Muñoz G, Gonzalo JA, Calleja C, González A, et al. The bispectral index scale: its use in the detection of brain death. *Transplant Proc.* 2005;37(9):3661-3663. doi:10.1016/j.transproceed.2005.08.054.
3. Hamal AQ, Bin Abdul Rehman AW. Artifact processing of epileptic EEG signals: an overview of different types of artifacts. In: 2013 International Conference on Advanced Computer Science Applications and Technologies. Piscataway (NJ): IEEE; 2013: 358-361.
4. Myles PS, Cairo S. Artifact in the bispectral index in a patient with severe ischemic brain injury. *Anesth Analg.* 2004;98:706-707. doi:10.1213/01.ANE.0000100780.04935.99.
5. Greer DM, Shemie SD, Lewis A, Torrance S, Varelas P, Goldenberg FD, et al. Determination of brain death/death by neurologic criteria: the World Brain Death Project. *JAMA.* 2020;324(11):1078-97. doi:10.1001/jama.2020.11586.
6. Saposnik G, Basile VS, Young GB. Movements in brain death: a systematic review. *Can J Neurol Sci.* 2009;36(2):154-60. doi:10.1017/S0317167100006516.
7. Myles PS, Leslie K, McNeil J, Forbes A, Chan MTV. Bispectral index monitoring to prevent awareness during anaesthesia: the B-Aware randomised controlled trial. *Lancet.* 2004;363(9423):1757-1763. doi:10.1016/S0140-6736(04)16287-9.
8. Bruhn J, Bouillon TW, Shafer SL. Bispectral index (BIS) and electromyographic activity. *Anesthesiology.* 2000;92(3):713-714.